

SUPREP BOWEL PREP

5 DAYS BEFORE PROCEDURE

- Stop all NSAIDs (Ibuprofen, Motrin, Advil, Naproxen, Aleve, Naprosyn, Mobic, Meloxicam, Diclofenac, and Celebrex). Tylenol and Aspirin 81mg are okay.
- Stop iron pills (Ferrous Sulfate, Fermalox, and Niferex)
- Stop foods that contain skins, hulls, seeds and nuts (peanuts, popcorn, grapes, beans, peels of apples/potatoes, strawberries, tomatoes, bread with seeds, rice, etc.). No salads.

1 DAY BEFORE PROCEDURE

- No solid foods
- No dairy products
- No red, orange, or purple liquids (Jell-O, sports drinks, juices or popsicles). Other colors are okay.
- Start clear liquid diet. Examples include: juice (white grape, white cranberry, lemonade), clear broth, soda (7-UP, Coke, etc.), black coffee and tea, popsicles, plain Jell-O (no fruit chunks), sports drinks, and Ensure Clear.

6PM the day before – drink first dose of Suprep solution

- Pour one 6 ounce bottle of Suprep liquid into the mixing container
- Add cool drinking water to the 16 ounce line on the container and mix
- Drink all of the liquid in the container
- You must drink 2 more 16 ounce containers of water over the next hour

DAY OF PROCEDURE

6 Hours before procedure

- Pour one 6 ounce bottle of Suprep liquid into the mixing container
- Add cool drinking water to the 16 ounce line on the container and mix
- Drink all the liquid in the container
- You must drink 2 more 16 ounce containers of water over the next hour

4 Hours before procedure

- **Stop** all waters/liquids
- **Stop** all candy, mints, chewing tobacco, smoking, and gum
- **Stop** diabetic medications but bring medications to procedure

MEDICATIONS

- Take heart, lung, blood pressure, and seizure medications with a sip of water the morning of procedure

MORNING OF PROCEDURE

- Take list of medications with you
- Wear comfortable clothes

Bring a responsible adult with you who will remain at the facility throughout your stay and will drive you home.

*****DO NOT EAT SOLID FOODS THE DAY BEFORE THE PROCEDURE. DO NOT EAT OR DRINK ANYTHING 4 HOURS PRIOR TO PROCEDURE TIME.*****

It is normal to feel abdominal cramping and bloating and have multiple loose stools during the prep period. Nausea and vomiting may also occur. Complete all of the prep if possible. A successful prep is when your bowel movements are a clear, yellow liquid.

Thompson

Colonoscopy Prep

**If you have any questions, call 812-743-5113 and ask for a nurse.